

## Bioinformatics 525: Module 2

### Introduction to Statistics

#### Lab #3

Read TROPHY.csv data in RStudio using “Import Dataset” on the Workspace Window.

**IMPORTANT:** type `attach(TROPHY)` to have the variables accessible for analysis.

1. **One-sample test:** Compare the proportion for a binary outcome to a given proportion value  $p_0$   
 $H_0: p = p_0$  vs.  $H_A: p \neq p_0$ .

- a. Create the following binary variables indicating a risk factor using a cutoff point. The “risk” category is coded as 1.

HDL01=1 if HDL  $\leq$  40

trig01=1 if Triglyceride  $\geq$  150

obese=1 if BMI24  $\geq$  30

- b. What are the number and the proportion of subjects with risk for the above variables?

- c. Visually display the number and the proportion of subjects with and without risk.

- d. Test if the proportion of subjects with HDL  $\leq$  40 is different from 40%:  $H_0: p = .4$  vs.  $H_A: p \neq .4$  using the chi-square test.

- e. Repeat d), but test  $H_0: p = .4$  vs.  $H_A: p \neq .4$  separately for males and females. Are the results the same?

2. **Two-sample test:** Compare the proportions of a binary outcome between two groups.

$H_0: p_1 = p_2$  vs.  $H_A: p_1 \neq p_2$ .

- a. Calculate the number and the % of males/females with HDL  $\leq 40$  and HDL  $> 40$

	Female	Male	Total
HDL $> 40$	n(col %)		
HDL $\leq 40$	p1	p2	p
Total	n1	n2	

- b. Visually display the number and the proportion of subjects with HDL  $\leq 40$  and HDL  $> 40$  for males and females. Is there any evidence of a difference by sex?

- c. Test whether the proportions of subjects with HDL  $\leq 40$  are different between males and females:  $H_0: p_1 = p_2$  vs.  $H_A: p_1 \neq p_2$  using one of the following tests:

i. Chi-square test

ii. Fisher's exact test

3. **Logistic Regression:** Identify metabolic risks of obesity (in each of the treatment group).
- Test if trig01 variable is a risk for obesity for subjects in the placebo group. Calculate the ARR, RR, and OR for obesity outcome by trig01 (< 150 vs. ≥ 150).
    - Use Chi-square test
    - Use Fisher's exact test
  - Repeat the same analysis as in a), but for subjects in the candesartan group. Are the findings from a) and b) the same? Does candesartan changes/modifies the effect of triglyceride on predicting obesity?
  - For the placebo group, use logistic regression model to predict obesity by trig01. Calculate the OR and compare it with the OR obtained from Fisher's exact test.

$$\log\left(\frac{\Pr(\text{Obese}=1)}{1-\Pr(\text{Obese}=1)}\right) = \beta_0 + \beta_1 * \text{trig01}$$

$$\text{OR}(\text{trig01}) = \exp(\beta_1)$$

- Repeat c), but use the following variables, trig01, insulin, and SBP0 to predict obesity. Calculate the OR for trig01 (i.e. adjusted OR). Compare the unadjusted OR from c) with the adjusted OR from d), are they the same?

$$\log\left(\frac{\Pr(\text{Obese}=1)}{1-\Pr(\text{Obese}=1)}\right) = \beta_0 + \beta_1 * \text{trig01} + \beta_2 * \text{Insulin} + \beta_3 * \text{SBP0}$$

$$\text{Adjusted OR for trig01: OR}(\text{trig01}) = \exp(\beta_1)$$

4. Please complete the muddy point assessment of this lab available at:  
<http://tinyurl.com/bioinf525-lab2-3>